

Local Anaesthetic

A03 Lite - Expires end of January 2022



This fact sheet is for general information about this procedure only. It is not intended to be used as medical advice or to replace advice that your relevant healthcare professional would give you. If you have a particular medical problem, please consult a healthcare professional.

© EIDO Systems International Limited. The operation and treatment information in this document is published under license from EIDO Systems International and is protected by copyright laws. Other than for your personal, non-commercial use, you may not copy, print out, download or otherwise reproduce any of the information.

eidohealthcare.com



What is a local anaesthetic?

A local anaesthetic is a drug that is injected into the tissues to make them numb. It can be used to give effective pain relief during and after an operation or procedure.

How does a local anaesthetic work?

Local anaesthetics temporarily stop nerves working so that you do not feel pain. The simplest form of local anaesthesia is to inject the anaesthetic just around the area where the operation is going to take place.

It is possible to numb all the nerves to an arm or a leg (a peripheral nerve block).

Local anaesthetics generally work for a few hours, depending on the type of anaesthetic and dose used. After this time the area should go back to normal.

Are there any alternatives to a local anaesthetic?

If you are concerned about being awake during the operation, it may be possible for it to be performed under a general anaesthetic.

Other forms of anaesthesia may be possible, such as an epidural or nerve block, although these also use local anaesthetics. Sometimes it may be possible to use a sedative as well as a local anaesthetic.

What complications can happen?

Some complications can be serious and can even cause death.

- Not enough pain relief
- Allergic reaction
- Bleeding
- Allergic reaction to the equipment, materials or medication
- Nerve damage
- Local anaesthetic toxicity

Summary

A local anaesthetic can be used for most people, usually giving a safe and effective form of pain relief during and after an operation or procedure.

Keep this information document. Use it to help you if you need to talk to the healthcare team.

Some information, such as risk and complication statistics, is taken from global studies and/or databases. Please ask your surgeon or doctor for more information about the risks that are specific to you.

This document is intended for information purposes only and should not replace advice that your relevant healthcare team would give you.

Acknowledgements

Reviewer: Iain Moppett (DM, MRCP, FRCA)